

PD-1 (M699V) Mouse mAb

CatalogNo: AH0001

•Main information

Target	PD1
Reactivity	Human
Applications	IHC,IF,ELISA
MW (kDa)	32kD (Calculated) 56kD (Observed)
Host Species	Mouse
Isotype	IgG1,Kappa

•Recommended Dilutions

IHC 1:200-400
IF 1:200-1:1000
ELISA 1:500-5000 Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0 (Cat#RH0011)

•Detailed information

Specificity	The antibody can specifically recognize human PD-1 protein.
Storage	-15°C to -25°C/1 year(Do not lower than -25°C, Ship by ice bag)
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Purification	The antibody was affinity-purified from ascites by affinity-chromatography using specific immunogen.
Clonality	Monoclonal
Clone Number	M699V

•Target Information

Gene name PDCD1 PD1

Protein Name Programmed cell death protein 1 (Protein PD-1) (hPD-1) (CD antigen CD279)


Organism	Gene ID	UniProt ID
Human	5133 ;	Q15116 ;

Cellular Localization Membranous, Cytoplasmic


Tissue specificity Placenta,Pooled tissue,Uterine cervix,

Function developmental stage:Induced at programmed cell death.,Disease:Genetic variation in PDCD1 is associated with susceptibility to systemic lupus erythematosus type 2 (SLEB2) [MIM:605218]. Systemic lupus erythematosus is a chronic, inflammatory and often febrile multisystemic disorder of connective tissue. It affects principally the skin, joints, kidneys and serosal membranes. It is thought to represent a failure of the regulatory mechanisms of the autoimmune system.,Function:Possible cell death inducer, in association with other factors.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:Monomer.,


•Validation Data




Human lymphoma tissue was stained with Anti-PD-1 Antibody




Human lymphoma tissue was stained with Anti-PD-1 Antibody



Human tonsil tissue was stained with Anti-PD-1 Antibody



Fluorescence multiplex immunohistochemical analysis of normal human appendix tissue (formalin-fixed paraffin-embedded section). The section was incubated in 3 rounds of staining; in the order of CK PAN., PD-1., Caldesmon pan. ,each using a separate fluorescent tyramide signal amplification system : Treble-Fluorescence immunohistochemical mouse/rabbit kit (pH9.0)



Zou, Ming-Xiang, et al. "A four-factor immune risk score signature predicts the clinical outcome of patients with spinal chordoma." Clinical and translational medicine 10.1 (2020): 224-237.

•Contact information

Orders: order@lamarck.cn
Support: support@lamarck.cn
Telephone: 400-801-6722
Website: <http://www.Lamarck.com>
Address: All rights reserved ©2025 Lamarck.



Please scan the QR code
to access additional
product information:
PD-1 (M699V)
Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)